

PAIRS OF FIXED POINTS FOR A CLASS OF OPERATORS ON HILBERT SPACES

ABDELHAK MOKHTARI^{a,b}, KAMEL SAOUDI^{c,d}, DUŠAN D. REPOVŠ^{e,f,g}

^aMathematics Department, Faculty of Mathematics and Informatics, University of M'sila, Algeria.

^bLaboratory of Fixed Point Theory and Applications, Department of Mathematics, E.N.S. Kouba, Algiers, Algeria

E-mail: abdelhak.mokhtari@univ-msila.dz

^cCollege of Sciences at Dammam, University of Imam Abdulrahman Bin Faisal, 31441 Dammam, Kingdom of Saudi Arabia

^dBasic and Applied Scientific Research Center, Imam Abdulrahman Bin Faisal University, P.O. Box 1982, 31441, Dammam, Kingdom of Saudi Arabia

E-mail: kmsaoudi@iau.edu.sa

^eFaculty of Education, University of Ljubljana, 1000 Ljubljana, Slovenia.

^fFaculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana, Slovenia.

^gInstitute of Mathematics, Physics and Mechanics, 1000 Ljubljana, Slovenia

E-mail: dusan.repovs@guest.arnes.si (Corresponding author)

Abstract. In this paper, existence of pairs of solutions is obtained for compact potential operators on Hilbert spaces. An application to a second-order boundary value problem is also given as an illustration of our results.

Key Words and Phrases: Hilbert space, potential operator, genus, fixed point theorem, boundary value problem.

2020 Mathematics Subject Classification: 35J25, 47G40, 47H10

1. INTRODUCTION

There are many papers that study the existence of fixed points for different types of operators, among the most important of these, the potential operator (or gradient operator) which can be regarded as the Gâteaux derivative of a suitable functional. Potential operators arise in many steady-state phenomena in physical problems from quantum mechanics, e.g. the potential of the Hamiltonian operator in the Schrödinger equation, see [3, 24, 25].

In fact, the variational methods to study linear and nonlinear equations are fully based on potential operators (see [4, 5, 11, 12, 22]). In [11], the authors considered nonlinear mappings $\phi \in C^1(H, \mathbb{R})$ defined on a Hilbert space H , ordered by a cone P and such that ϕ satisfies the Palais-Smale condition and has the expression

$$\phi' = I - A. \tag{1}$$

When A satisfies some growth conditions, A was shown to have a fixed point. A combination of topological and variational methods were used and an application to a second-order dynamic equation was given.

In [12], the authors discussed the existence of fixed points for a class of nonlinear operators on Hilbert spaces with lattice structure, by a combination of variational and partial ordered methods and they gave an application to second-order ordinary differential equations. In [23], the authors proved fixed point theorems for a widely more generalized hybrid non-self mappings on a Hilbert space. Using these results, they were able to prove the Browder-Petryshyn fixed point theorem [6] for strict pseudo-contractive non-self mappings and also generalized the fixed point theorem from [17] to the super hybrid non-self mappings.

Motivated by these previous works, we are concerned in this paper with the existence of pairs of fixed points on a Hilbert space H for an odd compact potential operator $A : H \rightarrow H$, satisfying the following sublinear growth condition

$$(\mathcal{H}) \quad \text{There exists } \theta \in [0, 1) \text{ such that } \limsup_{\|u\| \rightarrow +\infty} \frac{\|Au\|}{\|u\|^\theta} < \infty.$$

When $\theta = 1$, this condition is known as the quasiboundedness of A . If

$$\limsup_{\|u\| \rightarrow +\infty} \frac{\|Au\|}{\|u\|^\theta} < 1,$$

then existence of the fixed point of A is guaranteed by an application of Rothe's theorem (see, e.g., [13]).

The proofs of our results are based on the critical point theory. In particular, we apply Clark's theorem on a functional associated with the operator A . We choose it in an appropriate way so that its critical points are the same as the fixed points of A , as shown by relation (1). This method guarantees that the critical points on the boundary of the ball, that is different from the origin 0_H , which assures that the associated fixed points are nontrivial, which Schauder's theorem does not guarantee (see the next section).

This method requires the existence of linear operator and a set of unit vectors satisfying some properties. Moreover, this method naturally leads to multiplicity results which are rather difficult to achieve with classical fixed point theorems. We point out that Clark's theorem is often applied to prove the existence and multiplicity of weak solutions of boundary problems (see, e.g., [10, 19, 20]), but we believe that our work is among the first to apply it in the fixed point theory. Our main existence theorems are applied to a Dirichlet boundary value problems associated to a second-order ordinary differential equation. This simple model was chosen only to illustrate the effectiveness of the new fixed point theorems.

Our first existence result reads as follows.

Theorem 1.1. *Let H be a Hilbert space and $A : H \rightarrow H$ an odd compact potential operator satisfying assumption (\mathcal{H}) . Assume that there exist a linear operator B_1 on H , $r_1 > 0$, and $e_1 \in H$, with $\|e_1\| = 1$, such that*

$$(\mathcal{H}1) \quad (B_1(e_1), e_1) > 1,$$

$$(\mathcal{H}2) \quad (A(sr_1e_1), r_1e_1) \geq sr_1^2(B_1(e_1), e_1), \text{ for all } s \in (0, 1).$$

Then operator A has a pair of fixed points in $H \setminus \{0\}$.

Our second existence result reads as follows.

Theorem 1.2. *Let H be a Hilbert space and $A : H \rightarrow H$ an odd compact potential operator satisfying assumption (\mathcal{H}) . Assume that there exists a linear self adjoint operator B_2 on H such that*

$(\mathcal{H}1)'$ *there exist $e_2, e_3 \in H$ with $\|e_i\| = 1$, $i = 2, 3$, and $(e_2, e_3) = 0$, satisfying*

$$\begin{cases} (B_2(e_i), e_i) > 1, & i = 2, 3, \\ (B_2(e_2), e_3)^2 - \left(1 - (B_2(e_2), e_2)\right)\left(1 - (B_2(e_3), e_3)\right) < 0, \end{cases}$$

$(\mathcal{H}2)'$ *there exists a constant $r_2 > 0$ satisfying*

$$(A(su), u) \geq (B_2(su), u), \text{ for all } u \in \partial\mathcal{B}(0, r_2) \cap \langle e_2, e_3 \rangle, \text{ for all } s \in (0, 1),$$

where $\partial\mathcal{B}(0, r_2)$ denotes the boundary of the ball $\mathcal{B}(0, r_2)$ and $\langle e_2, e_3 \rangle$ is the subspace spanned by the vectors e_2 and e_3 .

Then operator A has two pairs of fixed points in $H \setminus \{0\}$.

We complete the introduction by an outline of the paper. In Section 2, we collect the necessary preliminary material. In Section 3, we prove the first existence result (Theorem 1.1). In Section 4, we prove the second existence result (Theorem 1.2). Finally, in Section 5, we give an application to a second-order boundary value problem to illustrate our results.

2. PRELIMINARIES

Let E be a real Banach space and Σ the class of all closed subsets $F \subset E \setminus \{0\}$ that are symmetric with respect to the origin, i.e., $u \in F$ implies $-u \in F$.

Definition 2.1. *Let $F \in \Sigma$. The Krasnosel'skii genus $\gamma(F)$ is defined as the least positive integer n such that there is an odd mapping $\varphi \in C(F, \mathbb{R}^n \setminus \{0\})$. If such n does not exist, then we set $\gamma(F) = +\infty$. Moreover, by definition, $\gamma(\emptyset) = 0$.*

Next we shall present a result on the computation of the genus that will be used in this work.

Proposition 2.1. *(see [21]) Let $F \subset E$, Ω be a bounded neighborhood of 0 in \mathbb{R}^N , and assume that there exists an odd homeomorphism $h \in C(F, \partial\Omega)$. Then $\gamma(F) = N$.*

More details on the genus can be found in [1, 7, 16, 18].

Definition 2.2. *Let $J \in C^1(E, \mathbb{R})$. If any sequence $(u_n) \subset E$ for which $(J(u_n))$ is bounded and $J'(u_n) \rightarrow 0$, as $n \rightarrow +\infty$ in E' , possesses a convergent subsequence, then we say that J satisfies the Palais-Smale condition (denoted by (PS) condition).*

The following theorem, due to Clark [9], will be crucial in the proof of our existence results.

Theorem 2.1. *Let $J \in C^1(E, \mathbb{R})$ be a functional satisfying the (PS) condition. Assume further that:*

- (a) J is even and bounded from below,

(b) *there is a compact set $K \in \Sigma$ such that $\gamma(K) = k$ and $\sup_{x \in K} J(x) < J(0)$.*

Then J possesses at least k pairs of distinct critical points and their corresponding critical values are less than $J(0)$.

We point out that this result is a consequence of a basic multiplicity theorem involving an invariant functional under the action of a compact topological group (see [9, 21]).

Recall that a mapping is said to be compact if it maps bounded sets into relatively compact sets. An operator $A : E \rightarrow E'$ is called a potential operator, if there exists a Gâteaux differentiable functional $T : E \rightarrow \mathbb{R}$ such that $T'(x) = A(x)$, for every $x \in E$ (see [8, 14]), where E' refers to the topological dual of E . Due to Avez [2], we know that, for all $u \in E$,

$$T(u) = \int_0^1 (A(su), u)_{E', E} ds.$$

Here $(\cdot, \cdot)_{E', E}$ refers to the duality pairing between E and its topological dual E' .

In this paper, we shall assume that $(H, (\cdot, \cdot))$ is a Hilbert space with (\cdot, \cdot) denoting the scalar product on H and $\|\cdot\| = \sqrt{(\cdot, \cdot)}$. Let A be a compact operator satisfying (\mathcal{H}) . Then there exist positive constants c, b such that for all $u \in H$, we have

$$\|Au\| \leq c\|u\|^\theta + b.$$

Indeed, by (\mathcal{H}) there exists some $R > 0$ such that for all $u \in H$ with $\|u\| \geq R$, we have $\|Au\| \leq c\|u\|^\theta$. Let $\mathcal{B}(0, R) \subset H$ be the ball centered at the origin with radius R . Since A is a compact operator, it follows that $\overline{A(\mathcal{B}(0, R))}$ is compact, whence $A(\mathcal{B}(0, R))$ is bounded, i.e. there exists $b > 0$ such that

$$\|Au\| \leq b, \text{ for all } u \in \mathcal{B}(0, R).$$

Then Schauder's fixed point theorem applies and yields a solution lying in the ball, which is possibly the trivial fixed point. On the other hand, our Theorem 1.1 will provide existence of a nontrivial fixed point u . Since A is odd, it follows that $-u$ is also a fixed point.

3. PROOF OF THEOREM 1.1

Consider the functional $J : H \rightarrow \mathbb{R}$ defined by

$$J(u) = \frac{1}{2}\|u\|^2 - \int_0^1 (A(su), u) ds, \quad (2)$$

where $Tu = \int_0^1 (A(su), u) ds$. Let $X_1 = \text{span}\{e_1\} \subset H$ be the subspace spanned by the vector e_1 and consider the set

$$K = \{u \in X_1 : \|u\| = r_1\} = \{\lambda e_1 : \|\lambda e_1\| = r_1\} = \{-r_1 e_1, r_1 e_1\}.$$

Step 1. It is clear that K is symmetric. Since X_1 and \mathbb{R} are isomorphic and K and S^0 are homeomorphic. Proposition 2.1 implies that $\gamma(K) = 1$; here S^0 refers to

the unit sphere in \mathbb{R} . Using Hypotheses $(\mathcal{H}1)$ and $(\mathcal{H}2)$, we get

$$\begin{aligned} J(-r_1 e_1) &= J(r_1 e_1) = \frac{1}{2} r_1^2 - \int_0^1 (A(sr_1 e_1), r_1 e_1) ds \\ &\leq \frac{1}{2} r_1^2 - \frac{1}{2} r_1^2 (B_1(e_1), e_1) = \frac{1}{2} (1 - (B_1(e_1), e_1)) r_1^2 < 0. \end{aligned}$$

It follows that

$$\sup_K J(u) = \max_{u \in \{-r_1 e_1, r_1 e_1\}} J(u) < 0 = J(0).$$

Step 2. J is bounded from below. Indeed,

$$\begin{aligned} J(u) &\geq \frac{1}{2} \|u\|^2 - \int_0^1 \|A(su)\| \|u\| ds \\ &\geq \frac{1}{2} \|u\|^2 - \|u\| \int_0^1 (c \|su\|^\theta + b) ds \\ &\geq \frac{1}{2} \|u\|^2 - \frac{c}{\theta+1} \|u\|^{\theta+1} - b \|u\|, \end{aligned} \tag{3}$$

and our claim follows from the fact that $\theta \in [0, 1)$.

Step 3. First, notice that since A is a potential operator, there exists a Gâteaux differentiable functional $T : H \rightarrow \mathbb{R}$ such that $T' = A$. More precisely,

$$T(u) = \int_0^1 \langle A(su), u \rangle ds.$$

Hence $J(u) = \frac{1}{2} \|u\|^2 - Tu$ and $J'(u)v = (u, v) - (Au, v)$, for all $v \in H$, that is $J' = I - A$.

Moreover, J satisfies the (PS) condition. Indeed, let (u_n) be a sequence in H such that $J'(u_n) \rightarrow 0$ and $J(u_n)$ is bounded. From (3), we get that (u_n) is bounded from below in H . Since A is compact, there exists a subsequence $(u_{n_k}) \subset H$ such that $A(u_{n_k}) \rightarrow v \in H$. Therefore $u_{n_k} \rightarrow v$ in H for

$$\|u_{n_k} - v\| \leq \|u_{n_k} - A(u_{n_k})\| + \|A(u_{n_k}) - v\| = \|J'(u_{n_k})\| + \|A(u_{n_k}) - v\|,$$

and the right-hand side tends to 0, as $k \rightarrow +\infty$.

By Theorem 2.1, we conclude that J has a pair of nontrivial critical points which are fixed points for the operator A . This completes the proof of Theorem 1.1. \square

4. PROOF OF THEOREM 1.2

We argue as in the proof of Theorem 1.1. Let $X_2 = \text{span}\{e_2, e_3\}$ be the subspace of H spanned by the vectors e_2 and e_3 and consider the set:

$$K' = \{u \in X_2 : \|u\| = r_2\} = \{\alpha e_2 + \beta e_3 : \alpha^2 + \beta^2 = r_2^2\}.$$

Clearly, $K' \in \Sigma$ and the homeomorphism

$$h : K' \ni u = \alpha e_2 + \beta e_3 \mapsto \left(\frac{\alpha}{\sqrt{\alpha^2 + \beta^2}}, \frac{\beta}{\sqrt{\alpha^2 + \beta^2}} \right) \in S^1$$

is odd. Now, Proposition 2.1 guarantees that $\gamma(K') = 2$. Here, S^1 is the unit sphere in \mathbb{R}^2 .

For $u \in K'$, assumption $(\mathcal{H}1)'$ now yields

$$\begin{aligned}
J(u) &= J(\alpha e_2 + \beta e_3) = \frac{1}{2}r_2^2 - \int_0^1 (A(su), u) ds \\
&\leq \frac{1}{2}r_2^2 - \int_0^1 (B_2(su), u) ds = \frac{1}{2}r_2^2 - \frac{1}{2}(B_2(\alpha e_2 + \beta e_3), \alpha e_2 + \beta e_3) \\
&= \frac{1}{2}r_2^2 - \frac{1}{2}\alpha^2(B_2(e_2), e_2) - \frac{1}{2}\alpha\beta(B_2(e_2), e_3) - \frac{1}{2}\alpha\beta(B_2(e_3), e_2) - \frac{1}{2}\beta^2(B_2(e_3), e_3) \\
&= \frac{1}{2}(1 - (B_2(e_2), e_2))\alpha^2 + \frac{1}{2}(1 - (B_2(e_3), e_3))\beta^2 - \alpha\beta(B_2(e_2), e_3).
\end{aligned}$$

Since the right-hand side term is strictly negative, it follows by compactness of K' that $\sup_{K'} J(u) < 0 = J(0)$. The rest of the proof parallels that of Theorem 1.1. This completes the proof of Theorem 1.2. \square

Remark 4.1. *If we attempt to extend the preceding results to the existence of n pairs of fixed points, then we can consider the $(n-1)$ -dimensional subspace*

$$K_n = \{u \in X_n = \langle e_1, e_2, \dots, e_n \rangle, \|u\| = r\} = \{\alpha_1 e_1 + \alpha_2 e_2 + \dots + \alpha_n e_n, \sqrt{\sum_{i=1}^n \alpha_i^2} = r\}.$$

Clearly, K is symmetric. If we assume that

$$(A(su), u) \geq (B(su), u), \quad \text{for all } u \in K_n, s \in (0, 1),$$

where B a linear self-adjoint operator, the question amounts to finding a sufficient condition for $\sup_{K_n} J(u)$ to be negative.

Let $u \in K_n$. Then, since B is linear and selfadjoint,

$$\begin{aligned}
J(u) &= \frac{1}{2}r^2 - \int_0^1 (A(su), u) ds \leq \frac{1}{2}r^2 - \frac{1}{2}(B(\sum_{i=1}^n \alpha_i e_i), \sum_{i=1}^n \alpha_i e_i) \\
&= \frac{1}{2}r^2 - \frac{1}{2}\sum_{i=1}^n \left(\sum_{j=1}^n \alpha_i \alpha_j (Be_i, e_j) \right) \\
&= \frac{1}{2} \left[\underbrace{\alpha_1^2 + \alpha_2^2 + \dots + \alpha_n^2}_{r^2} - \sum_{i=1}^n \left(\sum_{j=1}^n \alpha_i \alpha_j (Be_i, e_j) \right) \right].
\end{aligned}$$

However, we do not know whether the term on the right-hand side is negative.

5. APPLICATIONS

Example 5.1. *Consider the following boundary value problem*

$$\begin{cases} -u''(t) = f(t, u(t)), & t \in [0, 1], \\ u(0) = u(1) = 0, \end{cases} \quad (4)$$

where $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function. Clearly, solutions of problem (4) can be obtained as fixed points of the mapping A defined on H_0^1 by

$$Au(t) = \int_0^1 G(t, s)f(s, u(s)) ds, \quad (5)$$

where

$$G(t, s) = \begin{cases} t(1-s), & t \leq s, \\ s(1-t), & s \leq t \end{cases}$$

and H_0^1 is the standard Hilbert space with scalar product $(u, v) = \int_0^1 u'(t)v'(t)dt$ and the corresponding norm $\|u\| = \left(\int_0^1 |u'(t)|^2 dt\right)^{1/2}$. Then A satisfies

$$\begin{cases} -(Au)''(t) = f(t, u(t)), \\ (Au)(0) = (Au)(1) = 0. \end{cases} \quad (6)$$

Remark 5.1. One can check that the operator $A : H_0^1 \rightarrow H_0^1$ is compact (see [4]).

Next, we shall give an application to a second-order boundary value problem.

Theorem 5.1. Assume that there are positive functions $a_i \in L^1(0, 1)$, $a_i > 0$, a.e. $t \in (0, 1)$ ($i = 2, 3$), and $a_1 \in L^\infty(0, 1)$, where $m = \operatorname{ess\,inf}_{(0,1)} a_1 > 0$ and

$M = \operatorname{ess\,sup}_{(0,1)} a_1$, satisfying the following conditions

- (D₁) there exists $r_1 \in (0, 1)$ such that $\frac{f(t, u)}{u} \geq a_1(t)$, for all $u \in [-r_1, r_1] \setminus \{0\}$,
- (D₂) $f(t, u) \leq a_2(t)|u|^\theta + a_3(t)$, for all $u \in \mathbb{R}$ and some $\theta \in [0, 1)$,
- (D₃) there exist $e_1, e_2 \in H_0^1$ with $(e_1, e_2) = 0$ and $\|e_1\| = \|e_2\| = 1$ such that $m|e_i|_{L^2} > 1$, $i = 1, 2$,
- (D₄) $M^2 + 2\pi^2 m < \pi^4 + m^2$.

If f satisfies (f₃), then problem (4) has two pairs of nontrivial solutions in H_0^1 .

Proof. Let

$$J(u) = \frac{1}{2}\|u\|^2 - \int_0^1 F(t, u)dt, \quad (7)$$

where $F(t, u) = \int_0^u f(t, v)dv$. Using (D₁) together with the Lebesgue dominated convergence theorem, we can prove that $J \in C^1(H_0^1, \mathbb{R})$ and that for all $u, v \in H_0^1$,

$$\begin{aligned} J'(u)(v) &= \int_0^1 u'v' dt - \int_0^1 f(t, u(t))v(t) dt = \int_0^1 u'v' dt + \int_0^1 (Au)''(t)v(t) dt \\ &= \int_0^1 (u'v' - (Au)'(t)v'(t)) dt = (u, v) - (Au, v). \end{aligned}$$

Hence A is a potential operator. Define

$$Bu(t) = \int_0^1 G(t, s)a_1(s)u(s) ds. \quad (8)$$

We shall apply Theorem 1.2. The functional J and operator B are as given by (7) and (8). By (D₃), for $i = 1, 2$, we have

$$\begin{aligned} (B(e_i), e_i) &= \int_0^1 (B(e_i))'(t)_i'(t) dt = - \int_0^1 (B(e_i))''(t)e_i(t) dt \\ &= \int_0^1 a_1(t)e_i^2(t) dt \geq m|e_i|_{L^2}^2 > 1. \end{aligned}$$

Using the Cauchy-Schwarz and the Poincaré inequalities together with assumption (D_4) , we get the estimates

$$\begin{aligned}
& (B(e_1), e_2)^2 - \left(1 - (B(e_1), e_1)\right) \left(1 - (B(e_2), e_2)\right) \\
&= \left(\int_0^1 (B(e_1))'(t) e_2'(t) dt\right)^2 - \left(1 - \int_0^1 (B(e_1))'(t) e_1'(t) dt\right) \left(1 - \int_0^1 (B(e_2))'(t) e_2'(t) dt\right) \\
&= \left(\int_0^1 (B(e_1))''(t) e_2(t) dt\right)^2 - \left(1 + \int_0^1 (B(e_1))''(t) e_1(t) dt\right) \left(1 + \int_0^1 (B(e_2))''(t) e_2(t) dt\right) \\
&= \left(\int_0^1 a_1(t) e_1(t) e_2(t) dt\right)^2 - \left(1 - \int_0^1 a_1(t) e_1^2(t) dt\right) \left(1 - \int_0^1 a_1(t) e_2^2(t) dt\right) \\
&\leq M^2 |e_1|_{L^2}^2 |e_2|_{L^2}^2 - (1 - m |e_1|_{L^2}^2) (1 - m |e_2|_{L^2}^2) \\
&\leq M^2 |e_1|_{L^2}^2 |e_2|_{L^2}^2 - 1 + m |e_2|_{L^2}^2 + m |e_1|_{L^2}^2 - m^2 |e_1|_{L^2}^2 |e_2|_{L^2}^2 \\
&\leq \frac{1}{\lambda_1^2} M^2 + \frac{2}{\lambda_1} m - 1 - \frac{1}{\lambda_1^2} m^2 = \frac{1}{\pi^4} M^2 + \frac{2}{\pi^2} m - 1 - \frac{1}{\pi^4} m^2 < 0,
\end{aligned}$$

where $\lambda_1 = \pi^2$ is the first eigenvalue of the linear Dirichlet problem:

$$\begin{cases} -u''(t) = \lambda u(t), & t \in [0, 1], \\ u(0) = u(1) = 0. \end{cases}$$

Let $u \in H_0^1$ with $\|u\| = r_1$ and $s \in (0, 1)$. Since

$$|u|_\infty \leq \|u\| = r_1,$$

i.e., $-r_1 \leq u \leq r_1$, we have, by (D_1) ,

$$\begin{aligned}
(A(su), u) - (B(su), u) &= -\int_0^1 (A(su))''(t) u(t) dt + \int_0^1 (B(su))''(t) u(t) dt \\
&= \int_0^1 \frac{f(t, su(t))}{u(t)} u^2(t) dt - \int_0^1 a_1(t) s r e^2(t) dt \\
&= \int_0^1 \left(\frac{f(t, su(t))}{u(t)} - a_1(t) \right) u^2(t) dt \geq 0.
\end{aligned}$$

We can now deduce that conditions $(\mathcal{H}_1)'$ and $(\mathcal{H}_2)'$ are satisfied. Regarding condition (\mathcal{H}) , it is easy to see that it follows from condition (D_2) . Indeed, since the embedding

$H_0^1 \hookrightarrow C$ is continuous, we have

$$\begin{aligned} \|Au\| &= \sup_{\|v\|_{H_0^1} \leq 1} |(Au, v)| = \sup_{\|v\|_{H_0^1} \leq 1} \left| \int_0^1 (Au)'(t)v'(t) dt \right| \\ &= \sup_{\|v\|_{H_0^1} \leq 1} \left| \int_0^1 -(Au)''(t)v(t) dt \right| \leq \sup_{\|v\|_{H_0^1} \leq 1} \int_0^1 |f(t, u(t))v(t)| dt \\ &\leq c \int_0^1 (a_2(t)|u(t)|^\theta + a_3(t)) dt \leq c\|u\|^\theta \int_0^1 a_2(t) dt + \int_0^1 a_3(t) dt. \end{aligned}$$

Letting $c_1 = \int_0^1 a_2(t) dt$ and $c_2 = \int_0^1 a_3(t) dt$, we obtain

$$\limsup_{\|u\| \rightarrow +\infty} \frac{\|Au\|}{\|u\|^\theta} < \infty,$$

hence (\mathcal{H}) is satisfied.

Finally, Theorem 1.2 guarantees that operator A has two pairs of fixed points in H_0^1 . As a consequence, problem (4) admits two pairs of nontrivial solutions in H_0^1 . This completes the proof of Theorem 5.1. \square

Example 5.2. For any $\theta \in [0, 1)$ and $r_1, t \in (0, 1)$, consider the following function

$$f(t, u) = \begin{cases} a_2(t)u^\theta, & \text{if } u \geq 0, \\ -a_2(t)(-u)^\theta, & \text{if } u < 0, \end{cases}$$

where $a_i, i \in \{1, 2, 3\}$, are defined as follows

- $a_1(t) = 1 + t$,
- $a_2(t) = r_1^{1-\theta} a_1(t)$,
- $a_3(t) = t$.

Then function f satisfies the hypotheses of Theorem 5.1.

Acknowledgements. Repovš was supported by the Slovenian Research Agency grants P1-0292, N1-0278, N1-0114, J1-4031, J1-4001, and N1-0083. We thank the anonymous referees for their constructive comments and suggestions.

REFERENCES

- [1] A. Ambrosetti, A. Malchiodi, *Nonlinear Analysis and Semilinear Elliptic Problems*, Cambridge University Press, 2007.
- [2] A. Avez, *Calcul Différentiel*, Masson, Paris, 1983.
- [3] A. Balmaseda, D. Lonigro, J.M. Pérez-Pardo, *On the Schrödinger Equation for Time-Dependent Hamiltonians with a Constant Form Domain*, *Mathematics*, **10** (2022), art. 218.
- [4] A. Boucenna, S. Djebali, T. Moussaoui, *Fixed point theorems for compact potential operators in Hilbert spaces*, *Fixed Point Theory*, **18** (2017), no. 2, 493-502.
- [5] M. Briki, T. Moussaoui, *Fixed point theorems on Hilbert spaces via weak Ekeland variational principle*, *Electron. J. Math. Anal. Appl.*, **4** (2016), no. 2, 266-271.
- [6] F.E. Browder, W.V. Petryshyn, *Construction of fixed points of nonlinear mappings in Hilbert spaces*, *J. Math. Anal. Appl.*, **20** (1967), 197-228.
- [7] A. Castro, *Metodos variacionales y analisis funcional no linear*, X. Colóquio Colombiano de Matematicas, 1980.

- [8] J. Chabrowski, *Variational Methods for Potential Operator Equations*, Vol. 24, De Gruyter Studies in Mathematics, 1997.
- [9] D.C. Clark, *A variant of the Lusternik-Schnirelman theory*, Indiana Univ. Math. J., **22** (1972), 65-74.
- [10] F.J.S.A. Correa, G.M. Figueiredo, *On a p -Kirchhoff equation via Krasnoselskii's genus*, Appl. Math. Lett., **22** (2009), 819-822.
- [11] Y. Cui, J. Sun, *Fixed point theorems for a class of nonlinear operators in Hilbert spaces and applications*, Positivity, **15** (2011), 455-464.
- [12] Y. Cui, J. Sun, *Fixed point theorems for a class of nonlinear operators in Hilbert spaces with lattice structure and application*, Fixed Point Theory Appl., **2013** (2013), art. 345.
- [13] A. Granas, J. Dugundji, *Fixed Point Theory*, Springer Monographs in Mathematics, Springer-Verlag, New York, 2003.
- [14] E.W.C. Groesen, *Variational Methods for Nonlinear Operator Equations*, Proceedings Colloquium on Nonlinear Analysis, Mathematisch Centrum, Amsterdam, (1976), 100-191.
- [15] M. Imdad, A. Javid, *A Note on a Fixed Point Theorem in Hilbert Spaces*, Thai J. Math., **3** (2005), no. 2, 219-221.
- [16] O. Kavian, *Introduction à la théorie des points critiques et applications aux problèmes elliptiques*, Mathématiques & Applications, 13. Springer-Verlag, Paris, 1993.
- [17] P. Kocourek, W. Takahashi, J.C. Yao, *Fixed point theorems and weak convergence theorems for generalized hybrid mappings in Hilbert spaces*, Taiwan. J. Math., **14** (2010), 2497-2511.
- [18] M.A. Krasnosel'skii, *Topological Methods in the Theory of Nonlinear Integral Equations*, Pergamon Press, Macmillan Co., New York 1964.
- [19] A. Mokhtari, T. Moussaoui, D. O'Regan, *Existence and multiplicity of solutions for a $p(x)$ -Kirchhoff type problem via variational techniques*, Arch. Math. (Brno), **51** (2015), no. 3 163-173.
- [20] A. Mokhtari, T. Moussaoui, D. O'Regan, *Multiplicity results for an impulsive boundary value problem of $p(t)$ -Kirchhoff type via critical point theory*, Opuscula Math., **36** (2016), no. 5, 631-649.
- [21] P.H. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, CBMS Regional Conference Series in Mathematics, 65, American Mathematical Society, Providence, RI, 1986.
- [22] B. Ricceri, *Fixed points of nonexpansive potential operators in Hilbert spaces*, Fixed Point Theory Appl., **2012** (2012), art. 123.
- [23] W. Takahashi, N.C. Wong, J.C. Yao, *Fixed point theorems for nonlinear non-self mappings in Hilbert spaces and applications*, Fixed Point Theory Appl., **2013** (2013), art. 116.
- [24] K. Yajima, *Existence of Solutions for Schrödinger Evolution Equations*, Commun. Math. Phys., **110** (1987), 415-426.
- [25] K. Yajima, *Nonlinear Schrodinger Equation with Time Dependent Potential*, Commun. Math. Sci., **9** (2011), 937-964.

Received: ; *Accepted:*